

## Title

Evolution of *Escherichia coli* to 42°C and Subsequent Genetic Engineering Reveals Adaptive Mechanisms and Novel Mutations

## Authors

Troy E. Sandberg<sup>1</sup>, Margit Pedersen<sup>2</sup>, Ryan A. LaCroix<sup>1</sup>, Ali Ebrahim<sup>1</sup>, Mads Bonde<sup>2</sup>, Markus J. Herrgard<sup>2</sup>, Bernhard O. Palsson<sup>1,2</sup>, Morten Sommer<sup>2,3</sup>, Adam M. Feist<sup>1,2</sup>

1 Department of Bioengineering, University of California, San Diego, CA 92093, USA.

2 Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, 2800 Lyngby, Denmark.

3 Department of Systems Biology, Technical University of Denmark, 2800 Lyngby, Denmark.

\* Corresponding author. AM Feist, Department of Bioengineering, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0412, USA.

Tel.: 1 858 534 9592; Fax: 1 858 822 3120; E-mail: afeist@ucsd.edu

## Abstract

Adaptive laboratory evolution (ALE) has emerged as a valuable method by which to investigate microbial adaptation to a desired environment. Here we performed ALE to 42°C of ten parallel populations of *Escherichia coli* K-12 MG1655 grown in glucose minimal media. Tightly controlled experimental conditions allowed selection based on exponential-phase growth rate, yielding strains that uniformly converged towards a similar phenotype along distinct genetic paths. Adapted strains possessed as few as 6 and as many as 55 mutations, and of the 144 genes that mutated in total, 14 arose independently across two or more strains. This mutational recurrence pointed to the key genetic targets underlying the evolved fitness increase. Genome engineering was used to introduce the novel ALE-acquired alleles in random combinations into the ancestral strain, and competition between these engineered strains reaffirmed the impact of the key mutations on the growth rate at 42°C. Interestingly, most of the identified key gene targets differed significantly from those found in similar temperature adaptation studies, highlighting the sensitivity of genetic evolution to experimental conditions and ancestral genotype. Additionally, transcriptomic analysis of the ancestral and evolved strains revealed a general trend for restoration of the global expression state back towards pre-heat stressed levels. This restorative effect was previously documented following evolution to metabolic perturbations, and thus may represent a general feature of ALE experiments. The widespread evolved expression shifts were enabled by a comparatively scant number of regulatory mutations, providing a net fitness benefit but causing suboptimal expression levels for certain genes, such as those governing flagellar formation, which then became targets for additional ameliorating mutations. Overall, the results of this study provide insight into the adaptation process and yield lessons important for the future implementation of ALE as a tool for scientific research and engineering.

## Introduction

Adaptive laboratory evolution, or ALE, has developed over the years into a potent tool for biological discovery and engineering (Dragosits and Mattanovich 2013). By exploiting the inherent competition at play between organisms and the natural accumulation of mutations within a microbial population, desired phenotypic traits can be selected for without requiring *a priori* knowledge on how the traits might arise. These adaptively evolved organisms can then be subjected to whole genome resequencing, uncovering the genetic changes that enabled their phenotypic alteration. Additional data types, such as transcriptomics or metabolic uptake and secretion rates, serve to characterize the evolved strains and how they diverged from their ancestor, a divergence which must be enabled by their altered genotype. This analysis shines light on the functionality of particular genes (Cheng, et al. 2014) and the dynamics of the evolution process (Wiser, et al. 2013), increasing the biological knowledge base. While serving as a method to perform basic scientific inquiry such as this, ALE can be an equally useful tool for applied research, pairing with synthetic and systems biology to aid in the engineering of strains (Portnoy, et al. 2011).

ALE experiments often examine adaptation following a perturbation, either metabolic (e.g., growth on alternate carbon sources (Ibarra, et al. 2002) or following knockout of metabolic genes (Charusanti, et al. 2010)) or stressful (e.g., exposure to osmotic stress (Stoebel, et al. 2009) or high ethanol concentrations (Goodarzi, et al. 2010)). However, the selective pressure guiding the adaptation can also be influenced in large part by the environment in which the strain is evolved. Evolutionary environments typically involve either batch culturing, wherein populations (often several in parallel) are serially propagated to new flasks with fresh growth medium at regular intervals, or chemostats, in which growth in a bioreactor allows for tight control of nutrient levels and other factors such as pH and oxygenation. In either case, 'fitness' ostensibly refers to a growth advantage, but this becomes more complicated by the existence of spatial or temporal inhomogeneities in the culturing environment that can lead to ecological niches. For example, in a chemostat bacteria can persist by adhering to the walls of the bioreactor (Rao and Rao 2004), while batch cultures that reach stationary phase before passage can spawn subpopulations optimized for different phases of growth (Rozen, et al. 2009). If the target of investigation is the method by which a cell will evolve to a particular perturbation, it can be desirable to confine 'fitness' to a single aspect, reducing the potentially confounding variables towards which a population might additionally be evolving. To this effect, batch culture serial propagation in mid-exponential phase ensures that selection occurs primarily for growth rate.

In this study we sought to examine the ALE process by adaptively evolving the wild-type mesophilic bacterium *Escherichia coli* K-12 MG1655, arguably the most highly studied and well-characterized microorganism, to constant exponential growth at a stressful elevated temperature in glucose minimal media. Ten parallel populations were evolved at 42°C using an automated system allowing for passage of batch cultures in mid-exponential phase multiple times a day, enabling many generations of growth in a relatively short time. While genes that mutate in parallel across independently evolved populations are often taken to be the likely causes for the fitness increase (Wood, et al. 2005), true causal determination would require knocking-in each mutation to the starting strain in all possible combinations and comparing the resultant fitness, which would capture the individual effects of each mutation as well as their epistatic interactions. However, this quickly becomes prohibitively time consuming in strains with > 3 mutations, thus a different tool with which to probe mutational causality would be advantageous. For this reason we examined multiplex automated genome engineering (MAGE) (Wang, et al. 2009) as a technique to supplement ALE experiments. After identifying the mutations occurring in the endpoints of this evolution study, MAGE was used to introduce these ALE-acquired alleles in a random fashion into the starting strain, allowing the combinatorial knock-in method to be somewhat mimicked. By competing the heterogeneous populations of genetically engineered strains against one another and determining the mutants that frequently emerged victorious, causal mutations were identified and compared with those inferred by mutational parallelism across the ALE populations.

Elevated temperature was selected as the perturbation of interest for several reasons. Firstly, to aid in the analysis of mutational causality it would be beneficial to have more than simply one or two frequent gene targets, and adaptation to a global stress provided a diverse set of genetic changes. Additionally, a previous study (Tenailon, et al. 2012) investigated evolution of a large number of replicates in a very similar environment, differing only moderately in ancestral strain and method by which the batch cultures were serially propagated. Comparison with this work allowed examination of the extent to which mutational parallelism persists across studies. Furthermore, examining temperature stress was desirable because two other studies, both involving metabolic perturbations (Fong, et al. 2005; Carroll and Marx 2013), have provided strong evidence that a large feature of evolutionary adaptation involves acclimatizing changes in gene expression back towards pre-perturbed levels. By evaluating the transcriptome of our temperature-evolved strains we sought to determine whether this trend extended to stress perturbations, which would indicate that it may be a general feature of evolution, irrespective of the nature of the perturbation.

## Results

### Evolution Process and the Endpoint Phenotypes

Ten independent populations, started from wild-type *E. coli* K-12 MG1655, were adaptively evolved in M9 minimal media supplemented with 4 grams/liter glucose at 42°C for approximately 45 days. Cultures were serially passed (~5 times per day) to flasks with fresh media once reaching a target optical density such that stationary phase was never reached and glucose concentrations were always in great excess (never dropping below 3 g/L). As mutations accrued and gained dominance within the separate populations, their fitness increased markedly relative to the ancestral strain. The populations followed different trajectories along the fitness landscape, arriving at final growth rates on average 1.45 ( $\pm$  0.06, standard deviation) fold higher than the starting point (fig. 1; raw fitness data shown in supplementary fig. S1). The populations underwent approximately 1500 generations of growth (supplementary fig. S2A), but because mutations occur predominantly due to DNA polymerase errors in genomic replication during cell division (Camps, et al. 2003), the cumulative number of cell divisions (CCD) serves as a more meaningful scale for the time coordinate of an ALE than do generations (Lee, et al. 2011). This metric accounts for the population subsampling inherent to serial passage of cultures (supplementary fig. S2B). The CCD reached by the independent populations at the conclusion of the experiment ranged from 4.5 – 7.1 x 10<sup>12</sup> with an average of 5.5 x 10<sup>12</sup>. With the exception of the outlier experiment #1, CCD and final population fitness were significantly correlated (Pearson's  $r$  = 0.93 or 0.58 excluding or including experiment #1, respectively). This correlation is lost when using generations as the time coordinate (Pearson's  $r$  = 0.35 or 0.06 excluding or including experiment #1, respectively).

Clones were isolated from each of the evolved population endpoints and subjected to further analysis to determine the physiological differences at 42°C between the wild-type and the evolved strains (table 1, fig. 2). Interestingly, the fitness (i.e., growth rate) increase of the isolated clones relative to the ancestor is noticeably lower than that of the populations (paired  $t$ -test,  $p$  < 10<sup>-7</sup>). Although the discrepancy could be due to population-level altruistic interactions (Lee, et al. 2010), it may result simply from the evolved populations being fully physiologically adapted to constant exponential growth after ~1500 generations under static experimental conditions. This adapted intracellular state is in contrast with the evolved clones, for which growth curves were started up from stationary phase overnight cultures, potentially resulting in suboptimal performance due to insufficient time for re-acclimation of their protein expression machinery to the exponential growth, an effect previously documented (Shachrai, et al.

2010). Indeed, additional growth rate tests on the fastest and slowest growing endpoints clones (strains 4 and 9, respectively) and the populations from which they were isolated support this hypothesis, and reveal that the clone and population growth rates are in fact in excellent agreement (supplementary fig. S3). Thus, the selected clones were assumed to be representative of the dominant phenotype and genotype within the population endpoints.

The evolved strains all displayed similar physiological changes. However, genome examination revealed that strains 2 & 3 and 6 & 8 likely shared a lineage at some point (see Mutational Analysis), thus strains 3 and 8 were omitted from the following statistics to ensure that only fully independently evolved phenotypes were considered (changing which two strains to omit did not significantly alter any values). On average the independently evolved strains increased their growth rate ( $\mu$ ) by  $0.15 \text{ hr}^{-1}$  (equivalent to a decrease of 7.7 minutes in doubling time), increased their glucose uptake rate (GUR) by  $4.0 \text{ mmol gDW}^{-1} \text{ hr}^{-1}$ , increased their acetate production rate (APR) by  $3.9 \text{ mmol gDW}^{-1} \text{ hr}^{-1}$ , and decreased in biomass yield by  $0.07 \text{ gDW gGlc}^{-1}$  ( $Y_{X/S,ss}$ ; calculated at steady state by dividing growth rate by GUR). Correlation plots between the pairwise combinations of these characteristics highlight the physiological divergence of the evolved strains from the wild-type, as well as the relation or lack thereof between specific traits (fig. 3). There is a strong negative correlation (Pearson's  $r = -0.94$ ) between biomass yield and GUR among the evolved strains and a notable positive correlation ( $r = 0.74$ ) between APR and GUR, both of which contribute to the negative correlation between biomass yield and APR ( $r = -0.76$ ). These results imply that the strains adopted a similar phenotypic change, but to a varying extent, of increased glucose uptake at the cost of increased acetate overflow metabolism (Vemuri, et al. 2006), utilizing this greater metabolic flux not to create biomass but rather to generate more energy. Notably, growth rate itself was not correlated with any of the other traits (no Pearson's  $r$  with a magnitude above 0.22). These physiological trends replicate what is observed upon evolution to glucose minimal media at  $37^\circ\text{C}$  (LaCroix, et al. 2014).

### Mutational Analysis

Whole genome resequencing was performed on isolated clones from each experiment in order to investigate the genetic basis underlying their phenotypic changes. A total of 161 unique *de novo* mutations relative to the starting strain were found across all 10 endpoints (supplementary dataset S1), with a number of these being shared among two or more of the strains, or occurring within the same genes. The emergence of several 'hypermutators,' a recurring feature of many ALE experiments (Sniegowski, et al. 1997), accounted for the majority of these unique mutations. Four of the ten strains

were hypermutators, possessing on average 40 mutations, while the remaining six ‘non-mutators’ had an average of 7 mutations. The resequencing results indicate that two of the four hypermutators likely resulted from unanticipated cross-mixing between the evolving populations, thus only two hypermutators can be said to have occurred independently: one likely due to a SNP in *mutL* (strains 2 and 3) (Ban and Yang 1998) and the other due to a SNP in *mutD*, also known as *dnaQ* (strains 6 and 8) (Echols, et al. 1983). Analysis of the non-mutator genotypes does not indicate that they suffered from similar occurrences of cross-mixing (see Materials and Methods).

The observed mutations in the clones isolated from each population were compared. Fourteen genes or intergenic regions were found to mutate in parallel across two or more of the evolved strains (discounting those mutations shared due to cross-mixing), which fell into three general functional categories: mutations affecting metabolism, regulation, or the cell envelope (table 2). Intergenic mutations were categorized based on their position relative to the genes (e.g., SNPs downstream of one gene and upstream of another likely change expression of the latter) and transcriptomic data obtained from RNA-seq analysis. The key mutations within each of the three functional categories are now described:

- 1) Metabolic Mutations: Only three metabolic genes were found to mutate in more than one strain, but two of these occurred in half or more of the strains. Foremost among all mutations, regardless of category, is an 82 bp deletion between *pyrE* and *rph* that occurred in every strain except the two *mutD* hypermutators. This mutation does not appear to be specifically temperature-related, having been observed in several other ALE studies on adaptation to lactate- or glucose-supplemented minimal media at 30°C or 37°C, respectively (Conrad, et al. 2009; Blank, et al. 2014), and is hypothesized to relieve a defect in pyrimidine biosynthesis present in the starting strain (Jensen 1993). The second most predominant metabolic mutation was in *pykF*, or pyruvate kinase I, which experienced one intergenic SNP (accompanied by a 2.7 fold down-regulation in gene expression) and two different nonsynonymous mutations that may cause PykF inactivation through premature truncation of the enzyme (K286\*) or alteration of a putative substrate binding residue (T278S) (UniProt 2014). As with *pyrE/rph* this is likely not a temperature-specific beneficial mutation, given that PykF inactivation is a recurring feature of *E. coli* adaptation to glucose minimal media, hypothesized to allow for increased glucose uptake by decreasing the metabolism of phosphoenolpyruvate to pyruvate (Woods, et al. 2006; Kishimoto, et al. 2010; Blank, et al. 2014).

- 2) Regulatory Mutations: Five of the fourteen recurring mutations were found in regulatory genes, with three of these occurring in half of the strains: *rpoC*, *rne*, and *ygaH/mprA*. RpoC is a subunit of the RNA polymerase complex, which, given its ability to function as a global regulator, is a frequent target of mutations in bacterial adaptation (Murphy and Cashel 2003; Klein-Marcuschamer, et al. 2009). Similar ALE-identified *rpoC* mutations have been found to increase the general metabolic efficiency of *E. coli* grown in minimal media (Conrad, et al. 2010; Cheng, et al. 2014) or are inferred to adapt it to higher temperatures (Tenailon, et al. 2012). The ribonuclease *rne* had 4 different SNPs, the most diverse set of mutations observed in any one gene in this study. Rne is an essential enzyme involved in rRNA and tRNA processing and, as a key component of the RNA degradosome, is the rate-limiting or sole degrader of many transcripts (Jain, et al. 2002). A previously discovered *rne* mutant was nonviable at 44°C and significantly defective in rRNA/tRNA processing as well as mRNA degradation, but could be rescued at the elevated temperature by SNPs in the N-terminal catalytic domain (in which all four ALE-acquired SNPs occur) that restored the rRNA/tRNA processing to wild-type levels but did not undo the 2-3 fold decrease in mRNA decay rate (Perwez, et al. 2008). Relevantly, in prokaryotes the stability of mRNA is directly correlated with the optimal growth temperature of the organism (Gu, et al. 2010), suggesting that adaptation strategies to increased temperature might include increasing the stability of mRNA transcripts. Taken together, this implies that the *rne* mutations found in this study may function to improve the enzyme's rRNA/tRNA processing capabilities at 42°C without likewise improving its endonuclease efficacy. It is also of note that a SNP in *hfq* was observed in two strains, and Hfq binding can prevent mRNA degradation by Rne (fig. 4A) (Folichon, et al. 2003). The three different intergenic *ygaH/mprA* SNPs all occurred between 10-14 nucleotides upstream of the *mprA* start codon, likely influencing translation efficiency by modulation of the ribosomal binding site (Ringquist, et al. 1992). MprA is a transcriptional repressor for a number of genes that code for multidrug resistance pumps (Rodionov, et al. 2001). Although drug resistance is not a factor in this evolution, increased expression of the pumps leads to altered membrane flux for a variety of compounds (Piddock 2006), thus this regulatory mutation could also perhaps be classified as cell envelope-related.
- 3) Cell Envelope Mutations: Despite no single cell envelope-related gene being mutated in more than 3 of the 10 endpoints, there was nevertheless a clear selective pressure on the cell envelope in general, with 6 of the 14 recurring mutations falling into this category. These range from phospholipid transporters to membrane proteins and channels to genes involved in the levels of cell



envelope components. Note that although *nagC* and *nagA* could be classified as regulatory and metabolic, respectively, their role in the recycling of cell wall peptidoglycan (fig. 4B) is responsible for their classification as cell envelope-related mutations (Plumbridge 2009). This category of mutations is a feature of ALEs in general (Vijayendran, et al. 2008; Conrad, et al. 2009) and in this particular study may help the envelope maintain its essential physical properties at the elevated temperatures of the experiment (Sinensky 1974).

Although a number of ALE studies examining adaptation to increased temperature have been performed to date (Riehle, et al. 2001; Kishimoto, et al. 2010; Blaby, et al. 2012; Tenaillon, et al. 2012), the one by Tenaillon *et al.* serves as an ideal point of mutational comparison given the significant similarity in culture environment between our two studies and the large number of independent lines they evolved, providing a statistically significant basis against which to compare the mutations observed herein. However, despite the similarity in evolution environment, the difference in mutational frequency is extremely pronounced. Of the fourteen genes in this study that mutated in two or more strains, only four were found to mutate in *any* of the 114 evolved lines sequenced by Tenaillon. Both *rne* (4 different SNPs across 5 strains) and *ygaH/mprA* (3 different SNPs across 5 strains) had a mutation in only 1 out of the 114 lines. Mutations in *rpoC* and those in or around *ilvL* were the only noticeably recurring feature: 5 out of 10 and 2 out of 10 *rpoC* and *ilvL* mutants, respectively, in the evolved strains of this study, vs. 21 out of 114 and 29 out of 114 identified in the Tenaillon strains. Even given the relatively small sample size in this study, to have greater than 20% mutational infiltration of 12 genes in one instance and less than 1% in another implies a substantial difference in evolutionary trajectory. Though the recurring mutations in this study (table 2) overlap poorly with those identified by Tenaillon, there is slightly more overlap between the functional units determined to be significant in their study (possessing 5 or more mutations across the 114 strains) and the mutations observed in this work: 9 of the 26 functional units share one or more mutated genes, despite most mutational overlap occurring in only a single one of our evolved strains (supplementary table S1).

### **Analysis of Mutational Causality Using MAGE**

Multiplex automated genome engineering (MAGE) (Wang, et al. 2009) was utilized to examine causality of the various mutations identified in the evolved strains. To limit the scale of the experiments, only those mutations found in the 6 non-mutator strains were selected as targets to introduce via recombineering. This yielded 31 distinct mutations between the strains, resulting in 29 unique oligos with which to perform recombineering (two of the mutations were intergenic insertion sequences,

infeasible for use in MAGE due to their size, while insertion sequences occurring within genes were assumed to function equivalently to knock-outs). These oligos were organized into seven distinct pools – six pools which contained only those oligos corresponding to the mutations found within each of the non-mutator evolved strains (strains 1, 4, 5, 7, 9, and 10), and one pool which contained all 29 oligos. Nine rounds of recombineering were performed on the starting strain for each of these 7 pools, after which serial growth and passage of the cultures occurred to enrich for those strains existing within the highly heterogeneous starting populations that were most fit for growth at 42°C. Multiple colonies were isolated from each of these enriched populations and subjected to whole genome resequencing (supplementary dataset S2).

The most frequently observed MAGE mutations (occurring in at least three of the enriched strains, with a frequency > 25%) are given in table 3. Frequency for a gene is defined as the number of resequenced strains which possessed a mutation divided by the number of resequenced strains which potentially could have possessed it given the pool of oligos used in recombineering. This recapitulates what was observed in the evolved ALE strains – there is excellent agreement between the most frequent MAGE and ALE mutations, keeping in mind that the use of only non-mutator strain mutations in MAGE explains the absence of *yfdI*, *nagC*, *secD*, *dinQ/arsR*, *ilvL/ilvX*, and *hns/tdk* (fig. 5). This leaves *proQ*, a regulator of the membrane transporter ProP, as the sole disagreement between tables 2 and 3, which mutated in only ALE strain 7 (a missense SNP, Q216P) and yet was found in 9 of 19 relevant MAGE strains. These results reinforce the conclusions drawn from examination of mutational parallelism across the independently evolved ALE strains – that the identified genes are likely key targets for adaptation to growth at 42°C in glucose minimal media.

On average, enriched strains possessed 4 mutated genes targeted by recombineering and 3 mutations in secondary locations throughout the genome. However, several of the targeted genes experienced mutations that differed from what would be expected given the design of the recombineering oligos. With the exception of one, all of these novel, ‘off-target’ mutations occurred within the 70 bp region introduced into the genome by allelic replacement and thus likely resulted from incorporation of mis-synthesized oligos containing erroneous bases, as has been observed with MAGE previously (Wang, et al. 2009). Nevertheless, examining these unintentional mutations that were able to fix in the post-enrichment populations provides insight into the nature of the causal genetic changes.

In the case of *pyrE/rph*, 26 new SNPs and indels were found, and in no instance did a strain possess some combination of *rph* or *pyrE/rph* mutations that did not include the introduction of a shifted *rph*

stop codon. Rph is naturally defective in the *E. coli* strain used in this study (Jensen 1993), so alterations to its coding sequence should be phenotypically neutral. These mutations support the mechanism put forth previously (Conrad, et al. 2009), whereby a frameshift that moves the *rph* stop codon closer to the *pyrE* attenuator loop allows for improved regulation of *pyrE* expression. If these *rph* frameshifts and nonsense SNPs all yield roughly equivalent outcomes, then this begs the question of why only the same 82 bp deletion was observed in the ALE strains. This disparity in variation can be explained by the different acquisition method of ALE vs. MAGE mutations. The 82 bp *pyrE/rph* deletion is flanked by two 10 bp repeats, thus its prevalence in the adaptively evolved strains may be due to the relative frequency of DNA polymerase slippage during DNA replication (Michel 2000; Conrad, et al. 2009), and once this fixes within the populations there is no longer a selective force for continued genetic alteration in the region. This mutational bias is decreased under MAGE conditions – even with only a small fraction of oligos with synthesis errors, a variety of mutations can be acquired across each round of recombineering before the populations are subjected to growth-based selection in the form of enrichment, negating the ease-of-acquisition benefit the 82 bp deletion has in gaining dominance.

Unlike with *pyrE/rph*, where most frameshifts will be beneficial and gene inactivation is not a concern, one would expect that in the other MAGE gene targets the introduction of mis-synthesized oligos would in most instances result in a fitness decrease by altering the amino acid sequence of a functioning protein. This fitness decrease would render the mutants unable to survive the enrichment process, yielding few off-target mutations within the relevant genes of the enriched MAGE strains. The data support this conclusion; other than the 26 new *pyrE/rph* mutations, only 8 off-target mutations were found across all other MAGE-targeted genes in the resequenced strains, with more than half of these resulting in no protein sequence alteration (one intergenic 1 bp deletion, four synonymous SNPs, and three missense SNPs). There is no reason that the oligos used for *pyrE/rph* should be more erroneous than those for any other gene, thus the difference in number of off-target mutations that survived enrichment highlights the relative specificity of the ALE-identified mutations for effecting a fitness increase. However, *pykF* experienced no off-targets despite its two separate oligo-introduced mutations both potentially leading to gene inactivation, either by altering a potential binding residue or truncating the protein. It may be that these mutations only decreased the enzymatic activity of PykF rather than completely eliminating it, and reduced PykF functionality happens to be more beneficial than the complete inactivation that would likely result from the introduction of random, mis-synthesized oligos. It should also be noted that one off-target mutation, a SNP in *rpoC* (N762S), fell outside of an oligo-targeted region and thus may have arisen independently.

The resequenced MAGE strains were subjected to growth rate tests to ensure that they were in fact adapted to growth at 42°C. The tests revealed two strains that, despite possessing mutations and being picked from enriched populations, did not grow faster than the starting wild-type strain. These two strains were precluded from the MAGE mutational frequency analysis, but nevertheless highlighted an interesting feature: all adapted strains possessed mutations in *pyrE/rph* and/or *rpoC*, and the two unadapted strains were the only ones to have neither.

### Transcriptomic Profiling via RNA-seq

RNA-seq was performed to examine the global shifts in gene expression resulting from the altered genotypes of the evolved strains as compared to the wild-type ancestral strain at 42°C. To complement this analysis, the expression shift of the wild-type strain when grown at 42°C vs. 37°C was also determined (supplementary dataset S3). Figure 6A shows a heat map of fold changes for the 1208 genes deemed to be significantly differentially expressed ( $q$ -value < 0.05 (Storey and Tibshirani 2003)) in the wild-type strain when grown at the higher temperature. The differentially expressed wild-type genes and their pattern of up- and down-regulation are in good agreement with a previous study examining differential expression in *E. coli* after growth at 43°C (Gunasekera, et al. 2008), including such features as up-regulation of heat shock proteins (e.g., ClpB, DnaK, GrpE, and GroL, among others) and sulfur metabolism genes (*cys* genes), and down-regulation of genes involved in flagellar synthesis (*fli* and *flg* genes) and putrescine catabolism (*puu* genes).

In most of these cases, and indeed as a general trend across many of the genes (fig. 6A), the mutations of the evolved strains served to reverse the heat-induced transcriptional shift, restoring the expression state back towards the levels of the wild-type at 37°C. On average, each evolved strain had restorative shifts for 73% of the 1208 genes, and principal component analysis provides an additional means of visualizing these evolved shifts in gene expression (supplementary fig. S4). Such restoration has been documented in two other instances; in one study examining evolution of *E. coli* onto lactate or glycerol as the sole carbon source as opposed to glucose (Fong, et al. 2005), and in another examining evolution of *Methylobacterium extorquens* following replacement of a native central metabolism reaction with a functionally analogous, heterologous pathway (Carroll and Marx 2013). In both cases there was widespread restoration of expression back to the wild-type levels, highlighted by the less common ‘reinforcement’ of a change (i.e., down- or up-regulated genes at the start of evolution becoming increasingly down- or up-regulated following evolution, respectively). This reinforcement, when occurring across multiple of the independently evolved strains, was taken as evidence for the reinforcing

shift being an important factor in their increased fitness. In this study, highly parallel (occurring in at least 8 of the 10 evolved strains) transcriptional reinforcement occurs in 101 genes, compared to 703 genes that experience highly parallel restorative shifts. It should be noted that expression levels can be influenced by growth rate-dependent effects (Gyaneshwar, et al. 2005), but this cannot explain the observed transcriptional restoration given that the growth rate of the wild-type strain at 42°C is higher than at 37°C (0.82 hr<sup>-1</sup> vs. 0.7 hr<sup>-1</sup>), both of which are lower than the evolved strains.

The reinforcing shifts in gene expression across the temperature-evolved strains were examined in more detail. Evidence was found both supporting and refuting the assumption that these reinforcements point to key mechanisms of adaptation. Supporting data include the expression shifts observed in *nagE*, which increased 1.6 fold in the wild-type at 42°C followed by a further increase between 15-71 fold in evolved strains 1, 2, 6, and 8, in stark contrast with the remaining strains which changed in expression by no more than 1.6 fold (fig. 6B). The strains exhibiting the strongly reinforcing increase in *nagE* expression are those possessing mutations in *nagC* and/or *nagA*, both determined to be significant gene targets in the adaptation to increased temperature (table 2). These mutations are likely responsible for the observed *nagE* shifts, given the genes' mutual involvement in peptidoglycan recycling (Plumbridge 2009). This finding is similar to the strongly reinforced up-regulation of *pntAB* observed by Carroll and Marx in the *M. extorquens* evolved strains, which additional analysis showed to be a major contributor to their fitness increase, and the discovery of mutations within this operon.

In contrast with *nagE*, expression shifts in *proP* serve as an example of a reinforcing shift seemingly acting against the best interest of the cells. Upon initial growth at 42°C the wild-type strain decreases in *proP* expression by 1.5 fold, a decrease which is reinforced uniformly across the ten evolved strains. However, strain 7 exhibits the smallest expression decrease (an insignificant -1.2 fold, compared to an average of -2.7 fold across the other nine strains) and is also the strain that acquired a SNP in *proQ*. Though not appearing in multiple of the ALE endpoint strains, the *proQ* mutation was found to be significant in the MAGE enrichment analysis and is likely responsible for the greater comparative *proP* expression, given that ProQ's only known function is as a regulator of ProP levels (Chaulk, et al. 2011). This implies that the reinforced change in expression of *proP* is actually detrimental to growth at increased temperatures, enough to allow for fixation of this mutation that combats the change and provides a fitness increase of suitable magnitude to repeatedly survive MAGE enrichment. However, although no epistatic interactions could be inferred from the MAGE results (the *proQ* mutation did not

appear solely in the presence of any other single mutation), the possibility cannot be discounted that different expression shifts in *proP*, or indeed any gene, could have different effects in different strains.

In the same way that the reinforced expression shifts may not all necessarily aid in the fitness of the cells, so too is this the case for the restorative shifts. The most noticeably counterintuitive restorative shift is the massive up-regulation of flagellar genes in the evolved strains following their initial down-regulation in the wild-type upon growth at 42°C (fig. 6C). In the well-mixed environment of the ALE experiment there is no evident need for motility, so increased expression of these genes should incur a significant energetic cost (Soutourina and Bertin 2003) while providing no apparent benefit. Indeed, previous ALE studies, regardless of culturing temperature or carbon source in the growth medium, have yielded strains with mutations that led to the down-regulation of flagellar genes (Cooper, et al. 2003; Fong, et al. 2005; LaCroix, et al. 2014). Interestingly, strains 6 and 8 (the two *mutD* hypermutators) appear to have somewhat mitigated this flagellar up-regulation in the same way that strain 7 did for *proP* down-regulation. The SNP 22 base pairs upstream of the *hns* start codon that strains 6 and 8 share is a potential candidate for the genetic change behind their outlying behavior, given *hns*'s part in the regulation of flagellar genes (Landini and Zehnder 2002). Taken together, the data suggest that the evolved expression shifts, whether restorative or reinforcing, can sometimes be actively detrimental instead of beneficial.

Other than the influence of the *nagA/nagC* and *proQ* mutations on transcription of *nagE* and *proP*, respectively, the recurring ALE mutations (table 2) have only subtle, if at all discernible, effects on expression levels relative to the other endpoint strains lacking these mutations. This indistinct mutational influence on transcription includes both the genes that mutated, as well as separate genes that might logically experience an expression shift as a result (e.g., a *proQ* mutation causing *proP* shifts). Even the intergenic mutations, which are more likely to cause expression shifts than are protein sequence alterations (Wang, et al. 2011), fail to noticeably distinguish themselves from the strains without the mutations, and can be inconsistent in their influence. For example, although the strain possessing a SNP 59 base pairs upstream of the *pykF* start codon has the greatest decrease in *pykF* expression (-2.7 fold; the second largest decrease being -2.2 fold), it fails to stand distinctly apart from the remaining strains (*t*-test,  $p > 0.16$ ). Similarly, although 8 of the evolved strains share the identical 82 bp *pyrE/rph* deletion that should influence *pyrE* expression (Conrad, et al. 2009), their fold changes for the gene ranged between -3 and +5. Isolating the transcriptional effects of any individual mutation is clearly complicated by the presence of other genetic changes within the evolved strains. Elucidation of

the expression shifts enabled by specific mutations would likely require the creation and transcriptomic characterization of single knock-in mutants.

## Discussion

In this study, adaptive laboratory evolution of 10 replicate cultures was performed under carefully controlled conditions to yield *E. coli* strains selected solely for faster exponential-phase growth on glucose minimal media at 42°C. These strains were physiologically characterized, transcriptomically profiled, and subjected to whole genome resequencing to determine the genetic basis of their adaptation. Selected mutations were knocked-in to the ancestral strain in random combinations via MAGE to gain further insight into the causal genetic changes behind the acquired fitness increase. It was found that (1) frequency-based assignment of causality to mutations is largely consistent across the naturally evolved and genetically manipulated strains; (2) the path of genetic adaptation is greatly influenced both by the genotype of the starting strain and the conditions under which the evolution is performed; and (3) a variety of mutations leads to the same general trend of phenotypic and transcriptional convergence among the evolved strains, highlighted by the occasional counterintuitive shift in gene expression.

Although the number and variety of mutations capable of conferring a growth advantage in any particular environment may be nearly limitless, those which strongly contribute to an improved phenotype have a much greater probability of fixing within the evolving populations than do detrimental, neutral, or slightly beneficial mutations. Thus, genes mutated in parallel across independently evolved cultures stand out as likely causes for the observed fitness increase, rather than simply being ‘hitchhiker’ mutations (Wood, et al. 2005). This frequency-based analysis pointed to 14 different genes, fewer than 10% of the 144 genes mutated across all strains, as the foremost genetic targets for adaptation to growth on glucose at 42°C. A subset of these 144 altered genes, encompassing the mutations acquired by the six strains with few ( $\leq 8$ ) genetic changes relative to the common ancestor, were genetically engineered into the starting strain in random combinations with the techniques of MAGE. Repeated culturing of the highly heterogeneous MAGE populations enriched for the most fit strains existing within them, and genome resequencing revealed the genetic changes possessed by these strains. It was found that frequency-based analysis of the mutated genes within the MAGE strains perfectly recapitulated the likewise-determined key genes of the ALE strains (at least, for those genes which fell within the subset used in the genetic engineering) and pointed to only one

additional key gene target that did not happen to mutate in parallel in the ALE. Additionally, the unintentional inclusion of oligos with synthesis errors in the MAGE allelic replacement (Wang, et al. 2009) highlighted the effects the evolved mutations had on the genes; many unintentional off-target mutations accumulated if fitness was positively or neutrally affected by inactivation of the gene in question, but off-targets were generally lacking or synonymous if the mutated gene likely retained some measure of functionality. While not apparent in this study, MAGE analysis would also serve to highlight the existence of positive epistasis between acquired mutations if they predominantly showed up together in the post-enrichment strains, implying that their combined fitness benefit was greater than either individually. MAGE thus presents itself as a useful tool for reaffirmation of mutational causality following an ALE experiment.

Though the excellent agreement in mutational frequency between ALE and MAGE strains implies that these recurring mutations would similarly appear in other *E. coli* temperature evolution experiments, comparison with the work of Tenaillon *et al.* revealed that for the most part this is untrue. Of the 14 recurring ALE mutations identified herein, ten experienced zero mutations across all 114 lines evolved by Tenaillon, while two mutated in only 1 of the 114 lines. To understand where this discrepancy arises, it is important to classify the ways in which the two studies differ. Firstly, in this study K-12 MG1655 is used as the ancestral strain, differing from the B REL1206 strain used by Tenaillon (a descendant of B REL606 evolved for 2,000 generations on glucose minimal media at 37°C (Barrick, et al. 2009)). This strain difference provides a clear explanation for several of the mutational discrepancies: the 82 bp *pyrE/rph* deletion widespread among the strains generated here is specific to MG1655, thought to relieve an inherent defect in pyrimidine biosynthesis (Conrad, et al. 2009), and Tenaillon does not observe *pykF* mutations because B REL1206 already contains an inactivating mutation in *pykF*, precluding it from further selective forces. The use of different starting strains also provides a possible explanation for the lack of *rne* mutants found by Tenaillon. Rph is naturally defective in MG1655 and is known to interact *in vivo* with Rne (Duran-Figueroa, et al. 2006), so it may be that certain detrimental effects resulting from elevated temperature do not manifest themselves when Rph and Rne are able to form functional assemblies, but the absence of Rph places pressure upon Rne to develop compensatory mutations for improvement of its functionality. On top of these mutational discrepancies that can be intuited, there are doubtless those for which explanations are not obvious; for example, several beneficial *rpoB* mutations identified in the Tenaillon study were actually detrimental to fitness, or potentially even lethal, when introduced into MG1655 (Rodriguez-Verdugo, et al. 2013).



In addition to the different starting strains, differences in experimental methodology are likely to have a significant effect. Although evolved at essentially identical temperatures (42°C vs. 42.2°C) and on the same carbon source, the environments under which evolution occurred were nevertheless quite different. Whereas in this study constant exponential phase growth was maintained and all nutrients were always in great excess, thus allowing for selection based solely on growth rate, the methodology of Tenaillon mimics that of the long-term evolution experiment (LTEE) (Lenski, et al. 1991) – the media contained only 25 mg/liter of glucose (160 times less than the 4 g/liter used here) and cultures were passed once per day, as a result spending the majority of their time in stationary phase. Significant differences in bacterial behavior arise during stationary phase (Navarro Llorens, et al. 2010), and the process of repeated feast and famine causes the selective force to be non-constant. This can give rise to unusual features, such as coexisting subpopulations that thrive in different phases of the daily cycle (Rozen, et al. 2009). Furthermore, the Tenaillon strains evolved under poorly oxygenated conditions, given that 10 mL cultures were kept in 15 mL tubes shaken at only 100 rpm. The large difference in hypermutator prevalence between our two studies is also indicative of the methodological influence on genomic changes; at least 20% of our strains became hypermutators (discounting those likely caused by population cross-mixing), while Tenaillon evolved 115 lines of which only one became a hypermutator, and was precluded from further analysis.

As for the carbon source availability, in this study mutations facilitating faster glucose uptake are selected for only to the extent that they enable faster growth, whereas in LTEE conditions faster uptake of the scarce glucose in and of itself provides a significant advantage. This may be why under LTEE conditions *pykF* is known to acquire an internal insertion sequence (Barrick, et al. 2009), almost certainly leading to full gene inactivation, but the *pykF* mutations identified herein potentially preserve some of the enzyme's functionality given the failure of off-target *pykF* mutations to accrue in the MAGE strains. PykF is one of two enzymes catalyzing conversion of PEP to pyruvate in glycolysis, and it is hypothesized that PykF inactivation enables faster glucose uptake by increasing intracellular concentrations of PEP, which can be used by the phosphotransferase system to drive glucose uptake (Woods, et al. 2006). In the glucose-rich conditions of this study, PykF impairment as opposed to inactivation could serve as a more beneficial alternative, balancing the pros of increased glucose uptake against the cons of a decrease in glycolytic flux capabilities. Another relevant comparison exists between this study and one with identical experimental conditions and starting strain, but with the culturing temperature set to

37°C instead of 42°C (LaCroix, et al. 2014). The most notable genetic feature of the lower-temperature ALE was the various *rpoB* SNPs acquired by every evolved strain, but *rpoB* did not mutate in any of our higher-temperature ALE strains. Recurring mutational agreements exist only for *pyrE/rph*, *hns/tdk*, and *pykF* (which similarly experienced no alterations causing clear-cut inactivation). Together these mutational comparisons with other studies highlight the significant influence of experimental conditions and starting genotype on the resulting genetic adaptation. Thus, mutational reproducibility largely cannot be expected if even slightly different strains are used, if experimental methodology varies, or if evolution to multiple factors is simultaneous instead of sequential (e.g., evolving on glucose at 42°C vs. evolving first on glucose at 37°C, and subsequently at 42°C).

Despite notable recurrence of multiple mutations across the evolved strains of this study, other than *pyrE/rph* no gene was mutated in more than half of the endpoints. In spite of these mostly distinct genotypes, convergence occurred toward the same physiological and transcriptional state. Growth, glucose uptake, and acetate production rates increased uniformly across all strains, while biomass yield decreased. Many genes were significantly differentially expressed in the wild-type upon an upshift from 37°C to 42°C, but following evolution at the elevated temperature more than 70% of these genes reverted their expression state back towards that of the 'unperturbed' wild-type at 37°C. Mimicking the uniform nature of the physiological changes, these expression shifts were generally highly parallel – over 700 genes shifted back towards the unperturbed state in 8 or more of the 10 evolved strains. Though a minority, highly parallel shifts that moved expression further away from the unperturbed state occurred for over 100 genes. These trends in frequency and direction of shifts are in good agreement with two previous studies, demonstrating a consistent method of adaptation whether for growth on new carbon sources (Fong, et al. 2005), acclimation after perturbation of a metabolic pathway (Carroll and Marx 2013), or adjustment to a global stress such as elevated temperature. Any changes in physiology and gene expression observed between the wild-type and evolved strains must arise from the genetic differences that developed, thus the lack of uniformly-occurring mutations means that the different genotypes acquired by each strain result in roughly the same overall state.

That large-scale alterations to the expression state can result from relatively few genetic changes points to the influence of regulatory mutations, a category into which many of the most frequently mutated genes fell, most notably *rpoC*, *rne*, and *ygaH/mprA*. At least one of these genes was mutated in every evolved strain, a prevalence matched or exceeded only by mutations in *pyrE/rph* and *pykF*. Interestingly,

these latter two genes were the only metabolically-related mutations to occur in more than 2 of the 10 evolved strains, and neither is thought to result specifically from the increased temperature of the evolution, as discussed above; temperature-specific adaptation for the most part seems to involve changes in gene regulation or the cell envelope. The regulatory mutations clearly have a net positive benefit on the cell given their frequent occurrence, but this does not mean that every one of the widespread expression levels altered as a result is likewise beneficial. It could be that some expression shifts caused by the regulatory mutations are actively detrimental to the cell, just less so than the cumulative benefit of the global change. These deleterious changes would then be targets for further adaptation that could alleviate their effects. Here, as in previous works (Hindre, et al. 2012), we find several pieces of evidence supporting this hypothesis, most strikingly in the massive and counterintuitive up-regulation of flagellar genes following evolution, a trend that is resisted by two strains that succeeded in acquiring a mutation to mitigate the likely detrimental up-regulation. Although overall fitness is always selected for, it seems that over the course of an evolution certain subsystems of the cell may experience transient fitness decreases before additional mutations can develop to undo them.

Overall, the results of this study yield lessons important for the future implementation of ALE as a tool for both biological discovery and engineering. Firstly, we have demonstrated the utility of MAGE as a method by which to probe mutational causality, diminishing the need for the often-laborious creation of single and combinatorial knock-ins. Such a method would prove particularly useful in ALE studies lacking in replicates with which to examine mutational parallelism, or when hypermutability has produced an overabundance of genetic changes that obscure the strongly causal mutations hidden among them. Secondly, by comparing the mutations identified herein with those found in related studies, we have highlighted the great extent to which mutational development is predicated upon the ancestral genotype and influenced by all aspects of the evolutionary environment. Rarely will two studies examine evolution of identical strains under identical experimental conditions, thus authors should be wary of basing their expectations on comparisons with similar works. For example, if a more heat-tolerant production strain for a particular biochemical were desired, knocking-in the mutations identified in this study would not necessarily provide a benefit in the dissimilar genetic background and culturing environment. Finally, the previously documented restoration of expression state towards wild-type levels following metabolic perturbations and subsequent evolution has been shown to extend to global stresses, namely elevated temperature, as the perturbation. Transcriptomic analysis of our evolved

strains demonstrated this, as well as revealing the apparent occurrence of likely transient, localized, detrimental changes in gene expression. In much the same way that overall entropy maximization can be achieved despite entropic decreases in certain components, as with a protein folding in solution (Harano and Kinoshita 2005), genetic subsystems can experience fitness decreases in the cellular pursuit of overall fitness maximization. Although given enough evolutionary time such sub-optimal components would be ameliorated by compensatory mutations, this nevertheless provides an avenue for rational design to improve on the evolution process. Genome-scale metabolic models are capable of making *a priori* predictions on genes necessary for optimal cellular growth in a particular environment (Bordbar, et al. 2014), so those genes predicted to be useless could be knocked out ahead of time, saving the cell from potentially energetically-wasteful expression. Starting with a 'pre-adapted' strain such as this would expedite the evolution process and eliminate the need for mutations solely to counteract superfluous expression.

## Materials and Methods

### Adaptive Evolution

An automated system was used to propagate the evolving populations over the course of the ALE and monitor their growth rates. Flasks filled with 25 mL of 4 g/L M9 minimal media were kept at 42°C through placement in a heat block and aerated by magnetic tumble stirrers at 1800 RPM. At the start of the experiment, a flask of the wild-type strain *E. coli* K-12 MG1655 (ATCC47076) was grown up to stationary phase in the same conditions and used to inoculate 10 independent flasks with 900  $\mu$ L of culture. As the bacteria grew, the automated system took optical density (OD) measurements at 600 nm for each flask at four timepoints, targeted to evenly span an OD range of 0.05 - 0.3 based on the most recently calculated growth rate and the starting OD of the flask. Growth rates were determined by taking the slope of a least-squares linear regression line fit to the logarithm of the OD measurements. Once reaching the target OD of 0.3, 10  $\mu$ L of culture was passed into a new flask, and in the even numbered experiments this passage volume was changed to 100  $\mu$ L after 20 days of evolution. At the OD of 0.3, glucose concentration only dropped from 4 g/L to  $\sim$ 3.5 g/L (determined by HPLC measurement of the cultures), so exponential growth in excess glucose conditions was constantly maintained.

Growth rates for each flask were discarded as untrustworthy if fewer than four OD points were sampled, if the points spanned a range of fewer than 0.2 or more than 0.4 OD units, or if the  $R^2$  correlation was below 0.99. To reduce noise, the data were smoothed by averaging each point with its five adjacent neighbors on either side after applying weights following a normal distribution ( $\sigma = 2$ ) centered on the point in question. The evolution trajectory curves were obtained by fitting a monotonically increasing piecewise cubic spline to the smoothed data. Fitting to the unsmoothed data resulted in negligible changes to the spline. The cumulative number of cell divisions (CCD) was calculated as outlined previously (Lee, et al. 2011).

Glucose M9 minimal media consisted of 4 g/L Glucose, 0.1mM  $\text{CaCl}_2$ , 2.0mM  $\text{MgSO}_4$ , 1X Trace Element Solution and 1X M9 salts. 4000X Trace element solution consisted of 27g/L  $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ , 2g/L  $\text{ZnCl}_2 \cdot 4\text{H}_2\text{O}$ , 2g/L  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ , 2g/L  $\text{NaMoO}_4 \cdot 2\text{H}_2\text{O}$ , 1g/L  $\text{CaCl}_2 \cdot \text{H}_2\text{O}$ , 1.3g/L  $\text{CuCl}_2 \cdot 6\text{H}_2\text{O}$ , 0.5g/L  $\text{H}_3\text{BO}_3$ , and Concentrated HCl dissolved in ddH<sub>2</sub>O and sterile filtered. 10X M9 Salts solution consisted of 68g/L  $\text{Na}_2\text{HPO}_4$  anhydrous, 30g/L  $\text{KH}_2\text{PO}_4$ , 5g/L NaCl, and 10g/L  $\text{NH}_4\text{Cl}$  dissolved ddH<sub>2</sub>O and autoclaved.

## Cross-mixing Analysis

The sequencing results of the evolved endpoint strains highlighted the fact that there was unintentional cross-mixing between the populations over the course of the ALE. Strains 6 (41 total mutations) and 8 (55 total mutations) shared 24 identical mutations which did not occur in any of the other isolated strains, including the SNP which caused a truncated form of MutD (DnaQ) likely responsible for the observed hypermutator phenotype (Echols, et al. 1983). Although less immediately apparent, it seems likely that strains 2 (34 total mutations) and 3 (30 total mutations) suffered from the same cross-mixing – they share 3 identical mutations (a 1,222 bp deletion, a 1 bp deletion, and a SNP) which don't occur in any of the other strains, including the *mutL* mutation which likely causes their hypermutator phenotype (Ban and Yang 1998). For these reasons, we did not consider the mutations shared only between strains 6 & 8 or 2 & 3 to have arisen independently when performing mutational recurrence analysis.

Given these occurrences of cross-mixing it was important to establish that the non-mutator strains did not likewise share a partially-evolved ancestor, which would complicate the determination of recurring mutations. If it were the case that all identical mutations resulted from cross-mixing and did not arise independently, then we would expect to see a number of identical mutations in two or more strains, but these strains would not share identical mutations with any other strains. This was decidedly not the case (supplementary fig. S5). For example, consider the identical *rpoC* mutation shared between strains 5, 7, and 10. If we posit that this is the result of the same *rpoC* mutant invading and fixing within all three populations, then it must be that the remaining shared mutations all occurred independently (*hfq* in strains 1 & 7, *ygaH/mprA* in strains 4 & 5, and *pykF* & *miaE* in strains 4, 5, and 7) given that the other strains do not share the *rpoC* mutation. Thus in the 'worst case' scenario, only one of these four sets of shared mutations can be explained away by cross-mixing.

## DNA Sequencing

After colonies were isolated and selected on LB agar plates, genomic DNA was extracted using Promega's Wizard DNA Purification Kit. The quality of DNA was assessed with UV absorbance ratios using a Nano drop. DNA was quantified using Qubit dsDNA High Sensitivity assay. Paired-end resequencing libraries were generated using a Nextera XT kit from Illumina (San Diego, CA) with 1 ng of input DNA total. Sequences were obtained using an Illumina Miseq with a PE500v2 kit. The breseq pipeline (Barrick, et al. 2009) version 0.22 with bowtie2 (Langmead and Salzberg 2012) was used to map

sequencing reads and identify mutations relative to the *E. Coli* K12 MG1655 genome (NCBI accession NC\_000913.2). All samples had an average mapped coverage of at least 25x.

### **RNA Sequencing**

RNA-sequencing data were generated under conditions of exponential-phase, aerobic growth in glucose M9 minimal media. Cells were washed with Qiagen RNA-protect Bacteria Reagent and pelleted for storage at -80°C prior to RNA extraction. Cell pellets were thawed and incubated with Readylyse Lysozyme, SuperaseIn, Protease K, and 20% SDS for 20 minutes at 37°C. Total RNA was isolated and purified using the Qiagen RNeasy Mini Kit columns, following vendor procedures. An on-column DNase-treatment was performed for 30 minutes at room temperature. RNA was quantified using a Nano drop and quality assessed by running an RNA-nano chip on a bioanalyzer. Paired-end, strand-specific RNA-seq was performed following a modified dUTP method (Latif, et al. 2013). The rRNA was isolated using Epicentre's Ribo-Zero rRNA removal kit for Gram Negative Bacteria. Sequences were run on an Illumina Miseq using a PE50v2 kit. Reads were mapped to the *E. coli* K12 Genome (NC\_000913.2) using bowtie2, allowing for up to 2 mismatches and enforcing paired-end constraints. Differentially expressed genes were determined by cuffdiff (Trapnell, et al. 2010) with upper-quartile normalization and setting a maximum false discovery rate of 0.05. Principal component analysis was performed on the RNA-seq FPKM values using the *pca* function in MATLAB.

### **Physiological Measurements**

Growth curves for selected clones were started from stationary phase overnight cultures and grown under conditions identical to the ALE experiment, but with sampling occurring every 20 minutes. At each sampling time the OD<sub>600</sub> was taken along with a small volume of the growing culture that was then filter-sterilized. The filtrate was injected into a high performance liquid chromatography (HPLC) column (Aminex HPX-87H Column #125-0140). Compound concentrations at each time point were determined by comparison to a standard curve of known concentrations, and were used to determine rates of glucose uptake and acetate secretion by the cells. No other compounds were detected in the filtrate.

### **MAGE**

For recombineering, cells containing the pMA7 plasmid (manuscript under preparation) were grown in 15 mL LB media, with shaking at 37°C to an OD<sub>600</sub> of 0.4. Recombineering was mediated by the lambda Red beta protein and induced through the ParaBAD promoter for 10 minutes by adding arabinose to a

final concentration of 0.2%. After induction, cells were placed on ice for at least 15 minutes before being harvested, washed, and finally resuspended in a total volume of 200  $\mu$ L ice-cold sterile water. 50  $\mu$ L cells were mixed with 5 pmol oligo of each oligo and electroporated in 0.1 mm gap cuvettes; 1.8 kV, 200  $\Omega$ , 25  $\mu$ F. Immediately after electroporation, 1 mL LB was added to the cells. Cells were transferred to a 50 mL Falcon tube to a total volume of 5 mL LB and grown for at least 3 hours at 37°C to allow full segregation of chromosomal DNA. A total of 9 consecutive rounds of recombineering were performed. A total number of 29 oligos were designed (supplementary table S2) to recreate the majority of the mutations identified in ALE lines 1, 4, 5, 7, 9 and 10. Eight recombineering experiments were performed in parallel: six using the oligos corresponding to each of the non-mutator ALE lines, one using all 29 oligos, and one control using oligo P9 malk (table S2). Introduction of the malk oligo does not alter fitness in the glucose environments used, but makes the cells turn purple when grown on MacConkey agar maltose plates. In this control MAGE experiment, the recombinant formation frequency was found to be on average 10% by plating the recombineered population and determining the proportion of purple colonies. Recombinant formation frequency is influenced largely by oligo length (Wang, et al. 2009), which, along with all other experimental variables, was kept constant for all recombineering, thus allelic biases should not be present.


Following recombineering, the mixed cell populations were cultured and serially passed for three days, under the same conditions as the ALE experiment, to select for the strains most fit for growth at 42°C. These enriched MAGE populations were then streaked on minimal M9 glucose agar plates and single colonies were isolated, screened for improved growth rates at 42°C in a microtiter plate reader, and subjected to whole genome DNA resequencing.

## **Acknowledgements**


We thank Richard Szubin for his assistance with conducting experiments. This work was supported by the Novo Nordisk Foundation. TES is supported through the National Science Foundation Graduate Research Fellowship (grant DGE1144086).




## Figures




**Fig. 1. Fitness trajectories of the evolving populations.** Plotted is the population increase in fitness relative to the initial average growth rate of the common starting strain. At the approximate CCD of  $200 \times 10^{10}$ , the even-numbered experiments' serial passage volumes were increased 10 fold to examine the impact of CCD on overall fitness. The variability in CCD across experiments is due to this as well as fluctuations encountered in passage cell density over the course of the ALE. Dashed and dotted lines represent populations that became dominated by hypermutators, with identical colors indicating a similar hypermutator genotype.




**Fig. 2. Physiology of the temperature-evolved strains.** Strain numbers are listed below their respective bars. The error bars for growth rate represent standard deviation of three biological replicates, while error bars for the other traits represent 95% confidence intervals.




**Fig. 3. Pairwise correlation plots between the physiological traits of the evolved strains.** Strain numbers are listed next to their respective points, with the *mutL* hypermutators in yellow and the *mutD* hypermutators in red. The least squares linear best-fit line to the evolved strains (excluding the wild-type starting strain from the fit as well as strains 3 and 8, so that only fully independently evolved phenotypes are considered) is overlaid along with its Pearson correlation coefficient. ( $Y_{x/s,ss}$  = biomass yield at steady-state,  $\mu$  = growth rate, GUR = glucose uptake rate, APR = acetate production rate)



**Fig. 4. Repeatedly mutated genes with related functionality.** (A) Both *rne* and *hfq*, mutually involved in mRNA degradation, mutated in multiple of the temperature-evolved ALE strains. (B) Genes in the *nag* operon facilitate uptake of GlcNAc from the periplasmic space, which can be channeled away into glycolysis or reincorporated into the peptidoglycan of the cell wall. Genes in red (*nagC*, *nagA*) were recurring mutational targets. (OM = outer membrane, PG = peptidoglycan, IM = inner membrane, GlcNAc = N-acetylglucosamine, GlcNAc6p = N-acetylglucosamine-6-phosphate, GlcN6P = glucosamine-6-phosphate, F6P = fructose-6-phosphate)



**Fig. 5: Mutational agreement between the MAGE strain enrichment and ALE experiment results.** (A) A subset of the novel alleles acquired via evolution were randomly introduced into the ancestral strain with MAGE, and these MAGE strains were competed against one another in the enrichment process. 'Frequent ALE gene targets' refer to genes that mutated in at least two of the ten ALE endpoint strains, while 'Frequent MAGE gene targets' refer to those that mutated in at least 25% of the relevant post-enrichment MAGE strains. Circle areas are proportional to the number of genes contained within each category. (B) Frequency comparison between MAGE and ALE strains for the 9 most frequent MAGE gene targets. The only disagreement occurs for *proQ* (marked by \*), which was not a frequent ALE gene target. The vertical dashed line corresponds with 10%, the frequency for a gene mutating in only a single one of the ten ALE strains.



**Fig. 6. Heat maps of differential expression between the ancestral and evolved strains.** (A) The 1208 genes which are significantly differentially expressed ( $q$ -value  $< 0.05$ ) in the wild-type after growth at 42°C, rank-ordered. (B) Mutations acquired by the evolved strains (relevant strains marked with \*) could either serve to resist (*proP*; strain 7) or aid in (*nagE*; strains 1, 2, 6, and 8) the reinforcement of the WT expression shift. (C) Widespread up-regulation of flagellar genes after evolution, partially resisted by two strains (6 and 8) possessing similar genotypes.

## Tables

Table 1: Physiological characterization of colonies isolated from evolved population endpoints

Strain #	New Mutations Relative to WT	Growth Rate (hr <sup>-1</sup> )	Relative Fitness Increase	Glucose Uptake Rate (mmol gDW <sup>-1</sup> hr <sup>-1</sup> )	Acetate Production Rate (mmol gDW <sup>-1</sup> hr <sup>-1</sup> )	Biomass Yield (gDW gGlc <sup>-1</sup> )
WT	0	0.82±0.01	1	10.2±0.2	5.8±0.4	0.45±0.02
1	6	0.95±0.03	1.17±0.04	11.8±0.9	5.8±1.9	0.45±0.04
2 <sup>a</sup>	34	0.97±0.03	1.19±0.04	15.5±1.2	11.0±1.6	0.35±0.03
3 <sup>a</sup>	30	0.92±0.04	1.12±0.06	13.7±1.8	12.9±0.6	0.37±0.06
4	8	1.03±0.01	1.26±0.01	14.3±0.3	10.9±0.6	0.40±0.01
5	8	0.94±0.05	1.15±0.06	13.9±0.7	10.1±0.4	0.38±0.04
6 <sup>b</sup>	41	0.97±0.01	1.19±0.02	15.5±1.9	10.9±2.5	0.35±0.04
7	8	0.99±0.01	1.21±0.03	14.8±2.4	8.0±0.6	0.37±0.06
8 <sup>b</sup>	55	0.95±0.03	1.17±0.02	16.0±0.7	13.8±1.2	0.33±0.03
9	6	0.92±0.01	1.12±0.02	14.3±1.7	11.4±0.3	0.36±0.04
10	8	0.98±0.02	1.19±0.03	13.6±0.7	9.5±0.5	0.40±0.03

<sup>a</sup> and <sup>b</sup> are hypermutator strains of the same lineage. For growth rate and relative fitness the standard deviation based on triplicate experiments is given; for other values the 95% confidence interval is given

Table 2: Recurring mutations identified across the ALE endpoint strains

Gene	Specific Function	Category <sup>a</sup>	Mutation	Protein Change	Strain Number(s)
<i>pyrE/rph</i>	Orotate phosphoribosyl-transferase/RNase PH	M	82 bp deletion	frameshift	1, 2, 3, 4, 5, 7, 9, 10
<i>rpoC</i>	RNA polymerase subunit	R	SNP	A734V (GCG→GTG)	1, 2
			SNP	Q1367* (CAG→TAG)	5, 7, 10
<i>pykF</i>	Pyruvate kinase I	M	SNP	ydhZ/pykF intergenic (-498/-59)	3
			SNP	T278S (ACC→TCC)	10
			SNP	K286* (AAA→TAA)	4, 5, 7
<i>rne</i>	Ribonuclease E	R	SNP	D415N (GAC→AAC)	6/8
			SNP	H243Y (CAT→TAT)	3
			SNP	G124S (GGT→AGT)	9
<i>ygaH/mprA</i>	L-valine efflux transporter/MprA repressor	R	SNP	V19A (GTA→GCA)	2
			SNP	intergenic (+77/-14)	6/8
			SNP	intergenic (+80/-11)	10
<i>mleE</i>	Phospholipid ABC transporter	C	SNP	intergenic (+81/-10)	4, 5
			SNP	L107F (TTG→TTT)	4, 5, 7
			SNP	frameshift (519/1332 nt)	8
<i>yfdI</i>	Predicted inner membrane protein	C	SNP	Q186* (CAA→TAA)	6
			SNP	frameshift (1274/1332 nt)	3
<i>nagC</i>	PTS regulator	C	SNP	C264S (TGC→AGC)	6/8
			SNP	frameshift (218/1221 nt)	2
<i>hns/tdk</i>	H-NS regulator/thymidine kinase	R	SNP	intergenic (-22/-583)	6/8
			SNP	intergenic (-111/-486)	9
<i>hfq</i>	RNA binding protein	R	SNP	D9A (GAT→GCT)	1, 7
<i>nagA</i>	N-acetylglucosamine-6-phosphate deacetylase	C	SNP	G265D (GGC→GAC)	2
			SNP	In-frame (381-401/1149 nt)	1
<i>secD</i>	Membrane protein channel component	C	SNP	R181L (CGC→CTC)	8
			SNP	G499G (GGC→GGT)	2
<i>dinQ/arsR</i>	Toxic membrane peptide/metal-responsive regulator	C	SNP	intergenic (-209/-486)	8
			SNP	intergenic (-305/-390)	3
<i>ilvL/ilvX</i>	Leader peptide regulating isoleucine & valine biosynthesis operons	M	Double SNP	intergenic (+47/-39)	6
			SNP	intergenic (+48/-39)	8

<sup>a</sup> M, R, C = Metabolic, Regulatory, Cell envelope



Table 3: The most significantly recurring genes following enrichment of MAGE strains

<b>Gene</b>	<b>MAGE frequency</b>	<b>ALE frequency</b>	<b>New off-target mutations<sup>a</sup></b>
<i>pyrE/rph</i>	0.88	0.80	26 (including 9 frameshifts)
<i>ygaH/mprA</i>	0.86	0.50	1 (intergenic 1bp deletion)
<i>rpoC</i>	0.79	0.50	2 (I1357I, N762S)
<i>pykF</i>	0.55	0.50	0
<i>hfq</i>	0.50	0.20	0
<i>proQ</i>	0.47	0.10	1 (G212G)
<i>nagA</i>	0.36	0.20	2 (G127G, H129N)
<i>rne</i>	0.33	0.50	0
<i>mlaE</i>	0.27	0.30	1 (L99L)

<sup>a</sup> Protein alterations given in parentheses

## References

- Ban C, Yang W. 1998. Crystal structure and ATPase activity of MutL: implications for DNA repair and mutagenesis. *Cell* 95:541-552.
- Barrick JE, Yu DS, Yoon SH, Jeong H, Oh TK, Schneider D, Lenski RE, Kim JF. 2009. Genome evolution and adaptation in a long-term experiment with *Escherichia coli*. *Nature* 461:1243-1247.
- Blaby IK, Lyons BJ, Wroclawska-Hughes E, Phillips GC, Pyle TP, Chamberlin SG, Benner SA, Lyons TJ, Crecy-Lagard V, Crecy E. 2012. Experimental evolution of a facultative thermophile from a mesophilic ancestor. *Applied and environmental microbiology* 78:144-155.
- Blank D, Wolf L, Ackermann M, Silander OK. 2014. The predictability of molecular evolution during functional innovation. *Proceedings of the National Academy of Sciences of the United States of America*.
- Bordbar A, Monk JM, King ZA, Palsson BO. 2014. Constraint-based models predict metabolic and associated cellular functions. *Nature reviews. Genetics* 15:107-120.
- Camps M, Naukkarinen J, Johnson BP, Loeb LA. 2003. Targeted gene evolution in *Escherichia coli* using a highly error-prone DNA polymerase I. *Proceedings of the National Academy of Sciences of the United States of America* 100:9727-9732.
- Carroll SM, Marx CJ. 2013. Evolution after introduction of a novel metabolic pathway consistently leads to restoration of wild-type physiology. *PLoS genetics* 9:e1003427.
- Charusanti P, Conrad TM, Knight EM, Venkataraman K, Fong NL, Xie B, Gao Y, Palsson BO. 2010. Genetic basis of growth adaptation of *Escherichia coli* after deletion of *pgi*, a major metabolic gene. *PLoS genetics* 6:e1001186.
- Chaulk SG, Smith Frieday MN, Arthur DC, Culham DE, Edwards RA, Soo P, Frost LS, Keates RA, Glover JN, Wood JM. 2011. ProQ is an RNA chaperone that controls ProP levels in *Escherichia coli*. *Biochemistry* 50:3095-3106.
- Cheng KK, Lee BS, Masuda T, Ito T, Ikeda K, Hirayama A, Deng L, Dong J, Shimizu K, Soga T, et al. 2014. Global metabolic network reorganization by adaptive mutations allows fast growth of *Escherichia coli* on glycerol. *Nature communications* 5:3233.
- Conrad TM, Frazier M, Joyce AR, Cho BK, Knight EM, Lewis NE, Landick R, Palsson BO. 2010. RNA polymerase mutants found through adaptive evolution reprogram *Escherichia coli* for optimal growth in minimal media. *Proceedings of the National Academy of Sciences of the United States of America* 107:20500-20505.
- Conrad TM, Joyce AR, Applebee MK, Barrett CL, Xie B, Gao Y, Palsson BO. 2009. Whole-genome resequencing of *Escherichia coli* K-12 MG1655 undergoing short-term laboratory evolution in lactate minimal media reveals flexible selection of adaptive mutations. *Genome biology* 10:R118.
- Cooper TF, Rozen DE, Lenski RE. 2003. Parallel changes in gene expression after 20,000 generations of evolution in *Escherichiacoli*. *Proceedings of the National Academy of Sciences of the United States of America* 100:1072-1077.

Dragosits M, Mattanovich D. 2013. Adaptive laboratory evolution -- principles and applications for biotechnology. *Microbial cell factories* 12:64.

Duran-Figueroa NV, Pina-Escobedo A, Schroeder I, Simons RW, Garcia-Mena J. 2006. Polynucleotide phosphorylase interacts with ribonuclease E through a betabetaalphabetabetaalpha domain. *Biochimie* 88:725-735.

Echols H, Lu C, Burgers PM. 1983. Mutator strains of *Escherichia coli*, *mutD* and *dnaQ*, with defective exonucleolytic editing by DNA polymerase III holoenzyme. *Proceedings of the National Academy of Sciences of the United States of America* 80:2189-2192.

Folichon M, Arluison V, Pellegrini O, Huntzinger E, Regnier P, Hajnsdorf E. 2003. The poly(A) binding protein Hfq protects RNA from RNase E and exoribonucleolytic degradation. *Nucleic acids research* 31:7302-7310.

Fong SS, Joyce AR, Palsson BO. 2005. Parallel adaptive evolution cultures of *Escherichia coli* lead to convergent growth phenotypes with different gene expression states. *Genome research* 15:1365-1372.

Goodarzi H, Bennett BD, Amini S, Reaves ML, Hottes AK, Rabinowitz JD, Tavazoie S. 2010. Regulatory and metabolic rewiring during laboratory evolution of ethanol tolerance in *E. coli*. *Molecular systems biology* 6:378.

Gu W, Zhou T, Wilke CO. 2010. A universal trend of reduced mRNA stability near the translation-initiation site in prokaryotes and eukaryotes. *PLoS computational biology* 6:e1000664.

Gunasekera TS, Csonka LN, Paliy O. 2008. Genome-wide transcriptional responses of *Escherichia coli* K-12 to continuous osmotic and heat stresses. *Journal of bacteriology* 190:3712-3720.

Gyaneshwar P, Paliy O, McAuliffe J, Jones A, Jordan MI, Kustu S. 2005. Lessons from *Escherichia coli* genes similarly regulated in response to nitrogen and sulfur limitation. *Proceedings of the National Academy of Sciences of the United States of America* 102:3453-3458.

Harano Y, Kinoshita M. 2005. Translational-entropy gain of solvent upon protein folding. *Biophysical journal* 89:2701-2710.

Hindre T, Knibbe C, Beslon G, Schneider D. 2012. New insights into bacterial adaptation through in vivo and in silico experimental evolution. *Nature reviews. Microbiology* 10:352-365.

Ibarra RU, Edwards JS, Palsson BO. 2002. *Escherichia coli* K-12 undergoes adaptive evolution to achieve in silico predicted optimal growth. *Nature* 420:186-189.

Jain C, Deana A, Belasco JG. 2002. Consequences of RNase E scarcity in *Escherichia coli*. *Molecular microbiology* 43:1053-1064.

Jensen KF. 1993. The *Escherichia coli* K-12 "wild types" W3110 and MG1655 have an *rph* frameshift mutation that leads to pyrimidine starvation due to low *pyrE* expression levels. *Journal of bacteriology* 175:3401-3407.

Kishimoto T, Iijima L, Tatsumi M, Ono N, Oyake A, Hashimoto T, Matsuo M, Okubo M, Suzuki S, Mori K, et al. 2010. Transition from positive to neutral in mutation fixation along with continuing rising fitness in thermal adaptive evolution. *PLoS genetics* 6:e1001164.

Klein-Marcuschamer D, Santos CN, Yu H, Stephanopoulos G. 2009. Mutagenesis of the bacterial RNA polymerase alpha subunit for improvement of complex phenotypes. *Applied and environmental microbiology* 75:2705-2711.

LaCroix RA, Sandberg TE, O'Brien EJ, Utrilla J, Ebrahim A, Guzman GI, Szubin R, Palsson BO, Feist AM. 2014. Discovery of key mutations enabling rapid growth of *Escherichia coli* K-12 MG1655 on glucose minimal media using adaptive laboratory evolution. (under review).

Landini P, Zehnder AJ. 2002. The global regulatory *hns* gene negatively affects adhesion to solid surfaces by anaerobically grown *Escherichia coli* by modulating expression of flagellar genes and lipopolysaccharide production. *Journal of bacteriology* 184:1522-1529.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357-359.

Latif H, Lerman JA, Portnoy VA, Tarasova Y, Nagarajan H, Schrimpe-Rutledge AC, Smith RD, Adkins JN, Lee DH, Qiu Y, et al. 2013. The genome organization of *Thermotoga maritima* reflects its lifestyle. *PLoS Genet* 9:e1003485.

Lee DH, Feist AM, Barrett CL, Palsson BO. 2011. Cumulative number of cell divisions as a meaningful timescale for adaptive laboratory evolution of *Escherichia coli*. *PLoS one* 6:e26172.

Lee HH, Molla MN, Cantor CR, Collins JJ. 2010. Bacterial charity work leads to population-wide resistance. *Nature* 467:82-85.

Lenski RE, Rose MR, Simpson SC, Tadler SC. 1991. Long-Term Experimental Evolution in *Escherichia-Coli*. 1. Adaptation and Divergence during 2,000 Generations. *American Naturalist* 138:1315-1341.

Michel B. 2000. Replication fork arrest and DNA recombination. *Trends in biochemical sciences* 25:173-178.

Murphy H, Cashel M. 2003. Isolation of RNA polymerase suppressors of a (p)ppGpp deficiency. *Methods in enzymology* 371:596-601.

Navarro Llorens JM, Tormo A, Martinez-Garcia E. 2010. Stationary phase in gram-negative bacteria. *FEMS microbiology reviews* 34:476-495.

Perwez T, Hami D, Maples VF, Min Z, Wang BC, Kushner SR. 2008. Intragenic suppressors of temperature-sensitive *rne* mutations lead to the dissociation of RNase E activity on mRNA and tRNA substrates in *Escherichia coli*. *Nucleic acids research* 36:5306-5318.

Piddock LJ. 2006. Multidrug-resistance efflux pumps - not just for resistance. *Nature reviews. Microbiology* 4:629-636.

- Plumbridge J. 2009. An alternative route for recycling of N-acetylglucosamine from peptidoglycan involves the N-acetylglucosamine phosphotransferase system in *Escherichia coli*. *Journal of bacteriology* 191:5641-5647.
- Portnoy VA, Bezdán D, Zengler K. 2011. Adaptive laboratory evolution--harnessing the power of biology for metabolic engineering. *Current opinion in biotechnology* 22:590-594.
- Rao VSH, Rao PRS. 2004. Global stability in chemostat models involving time delays and wall growth. *Nonlinear Analysis-Real World Applications* 5:141-158.
- Riehle MM, Bennett AF, Long AD. 2001. Genetic architecture of thermal adaptation in *Escherichia coli*. *Proceedings of the National Academy of Sciences of the United States of America* 98:525-530.
- Ringquist S, Shinedling S, Barrick D, Green L, Binkley J, Stormo GD, Gold L. 1992. Translation initiation in *Escherichia coli*: sequences within the ribosome-binding site. *Molecular microbiology* 6:1219-1229.
- Rodionov DA, Gelfand MS, Mironov AA, Rakhmaninova AB. 2001. Comparative approach to analysis of regulation in complete genomes: multidrug resistance systems in gamma-proteobacteria. *Journal of molecular microbiology and biotechnology* 3:319-324.
- Rodriguez-Verdugo A, Gaut BS, Tenaille O. 2013. Evolution of *Escherichia coli* rifampicin resistance in an antibiotic-free environment during thermal stress. *BMC evolutionary biology* 13:50.
- Rozen DE, Philippe N, Arjan de Visser J, Lenski RE, Schneider D. 2009. Death and cannibalism in a seasonal environment facilitate bacterial coexistence. *Ecology letters* 12:34-44.
- Shachrai I, Zaslaver A, Alon U, Dekel E. 2010. Cost of unneeded proteins in *E. coli* is reduced after several generations in exponential growth. *Molecular cell* 38:758-767.
- Sinensky M. 1974. Homeoviscous adaptation--a homeostatic process that regulates the viscosity of membrane lipids in *Escherichia coli*. *Proceedings of the National Academy of Sciences of the United States of America* 71:522-525.
- Sniegowski PD, Gerrish PJ, Lenski RE. 1997. Evolution of high mutation rates in experimental populations of *E. coli*. *Nature* 387:703-705.
- Soutourina OA, Bertin PN. 2003. Regulation cascade of flagellar expression in Gram-negative bacteria. *FEMS microbiology reviews* 27:505-523.
- Stoebel DM, Hokamp K, Last MS, Dorman CJ. 2009. Compensatory evolution of gene regulation in response to stress by *Escherichia coli* lacking RpoS. *PLoS genetics* 5:e1000671.
- Storey JD, Tibshirani R. 2003. Statistical significance for genomewide studies. *Proceedings of the National Academy of Sciences of the United States of America* 100:9440-9445.
- Tenaille O, Rodriguez-Verdugo A, Gaut RL, McDonald P, Bennett AF, Long AD, Gaut BS. 2012. The molecular diversity of adaptive convergence. *Science* 335:457-461.

Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold BJ, Pachter L. 2010. Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat Biotechnol* 28:511-515.

UniProt. 2014. Activities at the Universal Protein Resource (UniProt). *Nucleic acids research* 42:D191-198.

Vemuri GN, Altman E, Sangurdekar DP, Khodursky AB, Eiteman MA. 2006. Overflow metabolism in *Escherichia coli* during steady-state growth: transcriptional regulation and effect of the redox ratio. *Applied and environmental microbiology* 72:3653-3661.

Vijayendran C, Barsch A, Friehs K, Niehaus K, Becker A, Flaschel E. 2008. Perceiving molecular evolution processes in *Escherichia coli* by comprehensive metabolite and gene expression profiling. *Genome biology* 9:R72.

Wang HH, Isaacs FJ, Carr PA, Sun ZZ, Xu G, Forest CR, Church GM. 2009. Programming cells by multiplex genome engineering and accelerated evolution. *Nature* 460:894-898.

Wang L, Wang FF, Qian W. 2011. Evolutionary rewiring and reprogramming of bacterial transcription regulation. *Journal of genetics and genomics* 38:279-288.

Wiser MJ, Ribeck N, Lenski RE. 2013. Long-term dynamics of adaptation in asexual populations. *Science* 342:1364-1367.

Wood TE, Burke JM, Rieseberg LH. 2005. Parallel genotypic adaptation: when evolution repeats itself. *Genetica* 123:157-170.

Woods R, Schneider D, Winkworth CL, Riley MA, Lenski RE. 2006. Tests of parallel molecular evolution in a long-term experiment with *Escherichia coli*. *Proceedings of the National Academy of Sciences of the United States of America* 103:9107-9112.